



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Snowy Owls were quite abundant during November and December, 1896, and four were taken in a steel trap set on the top of a long pole in an open field. Goshawks were also unusually abundant during the winter, and what is more unusual, were nearly all in adult plumage, and mostly males. Of twenty-one captured, nineteen were in adult plumage, and eighteen were males. We usually have the young birds in far greater numbers than the adult. Pine and Evening Grosbeak were present but not in great numbers. Lesser Redpolls and Snowflakes were also represented in small numbers.

J. N. CLARK, *Meridian, Wis.*

PINE SISKIN, *Spinus pinus*.—Further notes upon the unprecedented southward movement of this species have been received; this time from the extreme south. Writing of his visit in Louisiana, Mr. Widmann says: "I found it first March 2, a flock of fifty, near Mandeville, La., north of Lake Pontchartrain, and again March 5, at Madisonville, La., also a good sized flock in company with Goldfinches. Then again March 13, in the woods on the right bank of the Mississippi opposite New Orleans, a few, of which Mr. Kopman, a local ornithologist, took a specimen, the first he had ever seen. Prof. Beyer and Mr. Allison, the other ornithologists of New Orleans, confirm his statement, and consider it a rare find."

The species was fairly common at Oberlin, May 22.

AMERICAN CROW, *Corvus americanus*.—While out collecting Crow eggs on April 22, 1897, with Mr. F. L. Burns, I found a nest containing two young Crows several days old, and Mr. Burns took a set of six eggs from a nest lined with human hair. This is rather an early date for young.

RUSSELL GRAY, *Philadelphia, Pa.*

Is it generally known that early in the season the Towhee has a lengthy, fragmentary, soliloquizing song? The general effect of it is like the song of Brown Thrasher when heard in the distance. The Towhee utters this song as he works about the brush-piles, his scratching not interfering with the song. In scratching for food he hops forward and then kicks back with both feet simultaneously, alighting at his starting point. Beetles and other articles of food are thoroughly masticated before being swallowed.

J. C. GALLOWAY, *Montgomery, Ohio.*

WARBLING VIREO, *Vireo gilvus*.—During the collecting season of 1896, having occasion to collect a few sets of eggs and nests of three birds, eight sets were taken as follows: